




OVERALL-HWC RANK: 39

CHANGE: A 4





Rank Based On: Weighted sum of the number of standard deviations each measure is from the national average

STRENGTHS

- Low prevalence of smoking among women
- Low prevalence of tobacco use during pregnancy
- Low prevalence of tobacco use among adolescents

CHALLENGES

- Low prevalence of dental visits among women
- Low percentage of infants exclusively breastfed for six months
- High prevalence of 2+ adverse childhood experiences

HIGHLIGHTS

- Since the 2016 edition, smoking among women aged 18-44 decreased 13% from 13.5% to 11.7%
- Since the 2016 edition, the percentage of uninsured women aged 18-44 decreased 29% from 18.2% to 13.0%
- Since the 2016 edition, neonatal mortality decreased 5% from 3.7 to 3.5 deaths per 1,000 live births
- Since the 2016 edition, infant mortality increased 5% from 5.5 to 5.8 deaths per 1,000 live births
- Since the 2016 edition, tobacco use among adolescents aged 12-17 decreased 35% from 6.8% to 4.4%
- Since the 2016 edition, teen suicide increased 19% from 10.4 to 12.4 deaths per 100,000 adolescents aged 15-19

www.AmericasHealthRankings.org

32

33.0%



Arizona

Measure	Rating	2018 Value	2018 Rank
BEHAVIORS Chlamydia - Women (Cases per 100,000 females aged 15-44)	++	1,664.6	33
Excessive Drinking - Women (Percentage of women aged 18-44)		14.7%	7
Obesity - Women (Percentage of women aged 18-44)		25.7%	18
Physical Inactivity - Women (Percentage of women aged 18-44)	****	20.4%	20
Smoking - Women (Percentage of women aged 18-44)	*****	11.7%	3
Behaviors-Women * (Weighted z score)	+++++	0.030	8
Alcohol Consumption During Pregnancy † (Percentage of women with a recent live birth)	•	•	•
Breastfed (Percentage of infants)	+	15.6%	48
Sleep Position † (Percentage of women with a recent live birth)	•	•	•
Tobacco Use During Pregnancy (Percentage of live births)	+++++	5.3%	6
Behaviors-Infants* (Weighted z score)	+++	-0.034	30
High School Graduation (Percentage of students)	+	79.5%	43
Overweight or Obese-Youth (Percentage of children aged 10-17)	++++	26.9%	12
Substance Use Disorder-Youth (Percentage of children aged 12-17)	+	5.6%	42
Tobacco Use-Youth (Percentage of children aged 12-17)	*****	4.4%	4
Behaviors-Children* (Weighted z score)	+++	-0.006	22
Behaviors-HWC* (Weighted z score)		-0.010	21
POLICY (Parastance for each mat)		450/	40
Publicly-Funded Women's Health Services (Percentage of need met)		15%	46
Postpartum Visit † (Percentage of women with a recent live birth)			•
Uninsured Women (Percentage of women aged 18-44)		13.0%	34
Policy-Women* (Weighted z score)		-0.039	42
Baby-Friendly Facility (Percentage of live births)		2.2%	47
mPINC (Maternity Practices in Infant and Nutrition Care survey score)		79%	26
Policy-Infants* (Weighted z score)		-0.029	35
Adequate Health Insurance (Percentage of insured children aged 0-17)		73.3%	32
Children with Health Insurance (Percentage of children aged 0-17)		92.7%	41
Homeless Family Households (Number per 10,000 households)		4.2	20
Water Fluoridation (Percentage of population served by community water systems)		57.8%	38
Policy-Children* (Weighted z score)		-0.020	41
Policy-HWC* (Weighted z score) CLINICAL CARE	+	-0.087	42
Cervical Cancer Screening - Women (Percentage of women aged 21-44)	+	77.6%	42
Dedicated Health Care Provider - Women (Percentage of women aged 18-44)	+	66.8%	43
Dental Visit - Women (Percentage of women aged 18-44)	+	58.9%	48
Flu Vaccine - Women (Percentage of women aged 18-44)	+	28.6%	48
Well-Woman Visit (Percentage of women aged 18-44)	++	61.7%	38
Clinical Care-Women * (Weighted z score)	+	-0.071	47
Low-Risk Cesarean Delivery (Percentage of singleton, term, vertex, first births)		22.5%	14
Prenatal Care Before 3rd Trimester (Percentage of live births)	+	91.9%	42
Well-Baby Check (Percentage of babies aged 0-2)		91.3%	32
Clinical Care-Infants * (Weighted z score)		0.000	33
Adolescent Well-Visit (Percentage of children aged 12-17)		70.0%	48
Developmental Screening (Percentage of children aged 10 months-5 years)		26.4%	30
Immunizations - Adolescents (Mean z score)		-0.16	29
Immunization HPV Females (Percentage of females aged 13-17)		46.6%	31
Immunization HPV Males (Percentage of males aged 13-17)		41.7%	18
Immunization Meningococcal (Percentage of children aged 13-17)		85.2%	21
Immunization Tdap (Percentage of children aged 13-17)		84.3%	40
Immunizations - Children (Percentage of children aged 19-35 months)		69.9%	31
Medical Home for Child with Special Health Care Needs		38.4%	45
(Percentage of children aged 0-17 with special health care needs)			
Clinical Care-Children* (Weighted z score)	++	-0.023	37
Clinical Care-HWC* (Weighted z score)	+	-0.095	45
COMMUNITY & ENVIRONMENT Concentrated Disadvantage (Percentage of households)	+	33.8%	47
Food Insecurity - Household (Percentage of households)		14.6%	36
Intimate Partner Violence-Lifetime (Percentage of women aged 18 and older)		42.6%	47
Intimate Partner Violence-Pregnancy † (Percentage of women with a recent live birth)		42.U70 •	•
Community & Environment-Women* (Weighted z score)		-0.068	47
Household Smoke (Percentage of children aged 0-17)		14.8%	16
Infant Child Care Cost (Percentage of married-couple's median income)		13.9%	36
Community & Environment-Infants* (Weighted z score)		-0.009	23
Adverse Childhood Experiences (Percentage of children aged 0-17 with 2+ experiences)		30.6%	50
Adverse Childrood Experiences (Percentage of Children aged 0-17 with 2+ experiences)	7	30.6%	50

Neighborhood Amenities (Percentage of children aged 0-17) ++

Community & Environment-Children* (Weighted z score)	+	-0.055	44	
Community & Environment-HWC* (Weighted z score)	+	-0.131	47	
ALL DETERMINANTS			<u>-</u>	
All Determinants-Women * (Weighted z score)		-0.147	45	
All Determinants-Infants* (Weighted z score)		-0.072	39	
All Determinants-Children* (Weighted z score)		-0.104	45	
All Determinants-HWC* (Weighted z score)		-0.323	44	
OUTCOMES Diabetes - Women (Percentage of women aged 18-44)		2.3%	11	
Drug Deaths - Women (Deaths per 100,000 females aged 15-44)		15.7	26	
Intended Pregnancy † (Percentage of women with a recent live birth)		10.7		
Maternal Mortality (Deaths per 100,000 live births)		18.8	25	
Postpartum Depression † (Percentage of women with a recent live birth)		10.0	•	
Outcomes-Women * (Weighted z score)		0.017	19	
Infant Mortality (Deaths per 1,000 live births)		5.8	21	
Low Birthweight (Percentage of live births)				
		7.2%	17	
Neonatal Mortality (Deaths per 1,000 live births)		3.5	16	
Preterm Birth (Percentage of live births)		9.1%	19	
Outcomes-Infants* (Weighted z score)		0.029	17	
Child Mortality (Deaths per 100,000 children aged 1-18)		23.5	26	
Missed School Days (Percentage of children aged 6-17 missing 11+ days)		6.4%	48	
Teen Births (Births per 1,000 females aged 15-19)		26.3	36	
Teen Suicide (Deaths per 100,000 adolescents aged 15-19)		12.4	33	
Outcomes-Children* (Weighted z score)	++	-0.055	40	
Outcomes-HWC* (Weighted z score)	+++	-0.009	24	
OVERALL Overall-Women * (Weighted z score)		0.120	26	
, ,		-0.130 -0.043	36	
Overall-Infants* (Weighted z score)			31	
RATING RANK Overall-Children* (Weighted z score)		-0.159	44	
+++++ 1 - 10 Overall-HWC* (Weighted z score)	++	-0.332	39	
++++ 11 - 20				
* Value indicates z score. Negative scores are below US value; positive scores are above US value. For complete definitions of measures				

* Value indicates z score. Negative scores are below US value; positive scores are above US value. For complete definitions of measures including data sources and years, see "Data Sources and Measures".

† Not Included in Ranking

• Data Unavailable

++ 31 - 40

+ 41 – 50

Maternal Mortality

Teen Suicide



18.8



SINCE 2016, MATERNAL MORTALITY*
INCREASED 2.7% FROM 18.3 TO
18.8

* Deaths per 100,000 live births



12.4



SINCE 2016, TEEN SUICIDE*

INCREASED 19.2% FROM 10.4 TO 12.4

* Deaths per 100,000 adolescents aged 15-19

Child Mortality





23.5



SINCE 2016, CHILD MORTALITY*

DECREASED 5.6% FROM 24.9 TO
23.5

* Deaths per 100,000 children aged 1-18



15.7



SINCE 2016, DRUG DEATHS - WOMEN*
INCREASED 1.9% FROM 15.4 TO 15.7

* Deaths per 100,000 females aged 15-44

Breastfed

Tobacco Use During Pregnancy



15.6%

PERCENTAGE OF INFANTS
EXCLUSIVELY BREASTFED FOR SIX
MONTHS



5.3%



SINCE 2016, TOBACCO USE DURING PREGNANCY* **DECREASED 1.9% FROM 5.4% TO 5.3%**

* Percentage of live births



Top 5 States

1: MASSACHUSETTS 2: NEW HAMPSHIRE 3: RHODE ISLAND 4: VERMONT 5: MINNESOTA

AZ Value


120.0 150.0

120.0


150.0



Maternal Mortality





Teen Suicide



Page 5 of 8

Child Mortality





Race and ethnicity populations are as defined by the original source.

Page 6 of 8

AZ Value

65.0


65.0

52.0

52.0

Arizona

Drug Deaths - Women




Data suppression rules are as defined by the original source. Race and ethnicity populations are as defined by the original source.


Page 7 of 8



Trend: Breastfed, Arizona, United States



Trend: Tobacco Use During Pregnancy, Arizona, United States



Page 8 of 8